












Fluid dynamics: Life engine in the Ocean


Marina Lévy
LOCEAN-IPSL













Ocean mesoscale eddies




Sea-level Anomaly

Ocean mesoscale eddies





Atmospheric weather system - Clouds



Stirring of phytoplankton by ocean mesoscale eddies

Phytoplankton are invisible to the eye but visible from satellites






Gulf of Gascogne


100 km

Gulf of Finland








ESA Sentinel 2, 10 m resolution, Baltic Sea





Evolution over 8 days











z








Basin-scale patterns of phytoplankton

Ocean forests

Ocean deserts





Wind-driven vertical currents



Fishermen discovered fronts are hotspots



M.C. Escher, 1957




Prants, 2024

Seabirds hunt fronts too



Fregate birds foraging behavior in the Mozambique Channel

3D-GPS tracking + satellites





De Monte, 2012



Sea-surface temperature fronts

2018-02-15





Levy et al, 2018





REVIEW ARTICLE

DOI: [10.1038/s41467-018-07059-3](https://doi.org/10.1038/s41467-018-07059-3)

[OPEN](#)

The role of submesoscale currents in structuring marine ecosystems

Marina Lévy¹, Peter J.S. Franks² & K. Shafer Smith^{3,4}













REVIEW ARTICLE

DOI: [10.1038/s41467-018-07059-3](https://doi.org/10.1038/s41467-018-07059-3)

OPEN


The role of submesoscale currents in structuring marine ecosystems

Marina Lévy¹, Peter J.S. Franks² & K. Shafer Smith^{3,4}






Bridging scales: from one front to multiple fronts



Haeck et al., 2023



Bridging knowledges : from fronts to community structure




Annual Review of Marine Science

The Impact of Fine-Scale Currents on Biogeochemical Cycles in a Changing Ocean

Marina Lévy,¹ Damien Couespel,² Clément Haëck,¹
M.G. Keerthi,¹ Inès Mangolte,¹
and Channing J. Prend^{3,4}

Lévy et al, 2024



Lévy et al, 2025

$$\langle \overline{\partial_t C} \rangle = -\langle \nabla \cdot (\overline{C} \overline{\mathbf{v}}) \rangle - \langle \nabla \cdot (\overline{C'} \overline{\mathbf{v'}}) \rangle + \langle \partial_z (\overline{k_z} \partial_z \overline{C}) \rangle + \langle \partial_z (\overline{k'_z} \partial_z \overline{C'}) \rangle + \langle \overline{B_C} \rangle + \langle \overline{B'_C} \rangle$$

Mean
advection

Eddy
advection

Mean
Vertical
mixing

Eddy
Vertical
mixing

Mean
biogeochemical
reactions

Eddy
biogeochemical
reactions

$\langle . \rangle$: Average over bio-provinces and over periods > 1 year

$\langle \overline{\partial_t C} \rangle =$

0 : Mean state

> and < : Natural variability

long term trend: Climate change

$$0 = -\langle \nabla \cdot (\bar{C} \bar{\mathbf{v}}) \rangle - \langle \nabla \cdot (\bar{C}' \bar{\mathbf{v}}') \rangle + \langle \partial_z (\bar{k}_z \partial_z \bar{C}) \rangle + \langle \partial_z (\bar{k}'_z \partial_z \bar{C}') \rangle + \langle \bar{B}_C \rangle + \langle \bar{B}'_C \rangle$$

upscale connection

Mean
advection

Inverse
cascade

Finescales

Mean
Vertical
mixing


Direct
cascade

downscale
connection






Temperature change in France


Relative to average of 1961-2010 [°C]



Sea surface temperature trend







Biogeosciences, 17, 3439–3470, 2020
<https://doi.org/10.5194/bg-17-3439-2020>
 © Author(s) 2020. This work is distributed under
 the Creative Commons Attribution 4.0 License.




Biogeosciences
 Open Access
 EGU

Twenty-first century ocean warming, acidification, deoxygenation, and upper-ocean nutrient and primary production decline from CMIP6 model projections


Lester Kwiatkowski¹, Olivier Torres², Laurent Bopp², Olivier Aumont¹, Matthew Chamberlain³, James R. Christian³, John P. Dunne⁶, Marion Gehlen⁷, Tatiana Ilyina⁸, Jasmin G. John⁶, Andrew Lenton^{3,4}, Hongmei Li⁵, Nicole S. Lovenduski⁹, James C. Orr⁷, Julien Palmieri¹⁰, Yeray Santana-Falcón¹¹, Jörg Schwinger¹², Roland Séférian¹¹, Charles A. Stock³, Alessandro Tagliabue¹³, Yohei Takano^{8,14}, Jerry Tjiputra¹², Katsuya Toyama¹⁵, Hiroyuki Tsujino¹⁵, Michio Watanabe¹⁶, Akitomo Yamamoto¹⁶, Andrew Yool¹⁰, and Tilo Ziehn³




a) Decrease in PP



b) Deoxygenation



c) CO₂ uptake





Damping of climate change impacts and of climate change with finescales


due to upscale feedback, no change in eddy fluxes


Oceanic primary production decline halved in eddy-resolving simulations of global warming

Damien Couespel¹, Marina Lévy¹, and Laurent Bopp²

Sea surface temperature trend








Katia

Irma

Jose

NOAA, Septembre 2017



23 cm





Bleached coral reefs in Moorea, French Polynesia. **Alexis Rosenfeld / Getty Images**



Les centaines de milliers de poissons retrouvés échoués sur la plage de Quintana au Texas. - Quintana Beach Country Park








Photo: Alexis Rosenfeld





NOTES: Aquatic animals excluding aquatic mammals, crocodiles, alligators, caimans, aquatic products (corals, pearls, shells and sponges) and algae. Data expressed in live weight equivalent.

SOURCE: FAO. 2024. FishStat: Global production by production source 1950–2022. [Accessed on 29 March 2024]. In: FishStatJ. Available at: www.fao.org/fishery/en/statistics/software/fishstatj. Licence: CC-BY-4.0.



Photo: Alexis Rosenfeld








Photo: Alexis Rosenfeld



 SUSTAINABLE DEVELOPMENT GOALS





 The **sea is rising** – its level has increased by **23 cm** since 1901

 Global losses from **tropical storms and floods** hit **US\$ 102 billion**

 We have **set policies** to protect **8.34%** of the ocean

 **1,677** marine species are currently at **risk of extinction**

 Marine animal food production reached a **record** of **115 million tonnes**

 **37.7%** of fish stocks are **overexploited**

Starfishbarometer.org

Epilogue

